

The following is an index of the major topics addressed in the AANA Journal during 2018. The index is alphabetized according to headings, and entries appear in chronological order within the headings. Author and subject indexes are integrated.

- A**
- AANA Foundation: identifying patterns across the closed claim dataset using thematic analysis, 27; thematic analysis of obstetric anesthesia cases from the closed claims database, 464
- AANA Journal Courses: cryoneurolysis for the treatment of sensory nerve pain, 495; evidence-based use of nonopioid analgesics, 321; herbal medications and anesthesia case management, 242; hypoglossal nerve stimulation, 412; nitrous oxide for the management of labor analgesia, 72; residual neuromuscular blockade, 157
- Acceleromyography: implementation of, in increasing use of quantitative neuromuscular blockade monitoring, 269
- Adams JA: improved anesthesia handoff after implementation of the Written Handoff Anesthesia Tool (WHAT) [Lambert and Adams], 361
- Adams JS: isolated harlequin syndrome following brachial plexus nerve block via interscalene approach: a case report [Adams and Minzola], 56
- Adverse events: anesthetic management for happy puppet syndrome and, 67; during cosmetic surgery, 127; identifying patterns across the AANA Foundation closed claim dataset using thematic analysis, 27; improving patient outcomes through closed-claims analysis, 201
- Airway: securing following facial hematoma as complication in securing, 188
- Analgesia: evidence-based use of nonopioid analgesics, 321; intravenous dexamethasone as, 488; intravenous dexamethasone as an, 488; nitrous oxide for the management of labor, 72
- Anesthesia: bone cement implementation syndrome and, 433; case report of, for happy puppet syndrome, 67; in early recovery after surgical protocols for total knee and total hip arthroplasty, 32; evidence-based best practices of simulation, checklists, deliberate practice and debriefing in developing a regional training course, 119; evidence-based use of nonopioid analgesics, 321; fire risk assessment during time-out in the operating room, 99; herbal medications and, 242; implementation of standardized handoff of patients, 137; improved handoff after implementation of the Written Handoff Anesthesia Tool, 361; isolated harlequin syndrome following brachial plexus nerve block via interscalene approach, 56; ketamine in decreasing emergence delirium in surgical patients with posttraumatic stress disorder, 220; management of double-outlet right ventricle with Glenn shunt for Cesarean delivery, 408; maternal and fetal outcomes with early neuraxial engagement in obese parturients, 234; of patient with type 1 von Willebrand disease and uterine placental abruption, 209; spinal-epidural, for Cesarean delivery, 109; thematic analysis of cases from the AANA Foundation closed claims databank, 464; transfer of care in perioperative settings, 401; transfer of care in perioperative settings: a descriptive qualitative study [Wilbanks et al.], 401
- Anesthesiologist: converging or diverging concepts in patient management and decision making, online
- Angelman syndrome: anesthetic management for happy puppet syndrome, 67
- Antidepressant effects: investigation of Crocin in the male Sprague-Dawley rat, 225
- Antrum: point-of-care gastric ultrasound examination in management of ASA class 3E patient, 379
- Anxiolysis: investigation of Crocin in the male Sprague-Dawley rat, 225; nitrous oxide for the management of labor analgesia and, 72
- Apnea: in patients undergoing total joint arthroplasty, 282
- Aravindan A: anesthetic management of a unique case of double-outlet right ventricle with Glenn shunt for Cesarean delivery: a case report [Aravindan et al.], 408
- Arevalo VN: methylene blue as an adjunct to treat vasoplegia in patients undergoing cardiac surgery requiring cardiopulmonary bypass: a literature review [Arevalo and Bullerwell], 455
- Arthroplasty: anesthetic management in early recovery after surgery for total knee and total hip, 32; bone cement implementation syndrome and, 433
- Austin PN: anesthetic management of a patient with type 1 Von Willebrand disease and uterine placental abruption: a case report [Roberson et al.], 209; pharmacologic methods for preventing pruritus in patients receiving intrathecal opioids for Cesarean delivery [Vice-O'con et al.], 59
- Ayala MP: investigation of the anxiolytic and antidepressant effects of Crocin, a compound from Saffron (*Crocus sativus* L.), in the male Sprague-Dawley rat [Cerecuga et al.], 225
- B**
- Ball-valve obstruction: complications related to laryngeal mask airway use and, 443
- Bansal S: anesthetic management of a unique case of double-outlet right ventricle with Glenn shunt for Cesarean delivery: a case report [Aravindan et al.], 408
- B cells: effect of norepinephrine on intracellular Ca²⁺ levels in malignant hyperthermia, 383
- Beaumont DM: investigation of the anxiolytic and antidepressant effects of Crocin, a compound from Saffron (*Crocus sativus* L.), in the male Sprague-Dawley rat [Cerecuga et al.], 225
- Best practices: evidence based, of simulation, checklists, deliberate practice, and debriefing to develop a regional anesthesia training course, 119; intravenous fat emulsion for treatment of local anesthetic systemic toxicity, 290; for monitoring recovery from neuromuscular blockade, Supplement
- Blood disorders: anesthetic management of patient with type 1 von Willebrand disease, and uterine placental abruption, 209
- Bonds R: complications related to laryngeal mask airway use and a postoperative diagnosis of reinke edema: a case study [Cuevas et al.], 443; implementation of an education program for an ultrasound-guided liposomal bupivacaine transversus abdominis plane (TAP) block protocol for open

- abdominal procedures [O'Driscoll et al.], 479
- Bone cement implementation syndrome: understanding, 433
- Boudreau JA: nitrous oxide for the management of labor analgesia [Collins et al.], 72
- Boust R: a content and thematic analysis of closed claims resulting in death [Hranchook et al.], 311
- Boust RR: transfer of care in perioperative settings: a descriptive qualitative study [Wilbanks et al.], 401
- Brachial plexus: isolated harlequin syndrome following nerve block via interscalene approach, 56
- Bradley B: maternal and fetal outcomes with early neuraxial engagement in obese parturients [Ray and Bradley], 234
- Brain-derived neurotrophic factor (BDNF): effects of subanesthetic intravenous ketamine infusion on corticosterone in male Sprague-Dawley rats, 393
- Bramble RS: minimum effective dose (ED₅₀ and ED₉₅) of intrathecal hyperbaric bupivacaine for Cesarean delivery: a systematic review [Tubog et al.], 348
- Brown C: should continuous subcutaneous insulin infusion (CSII) pumps be used during the perioperative period? development of a clinical decision algorithm [Rotruck et al.], 194; understanding motivations and barriers of CRNAs involved in global health: a qualitative descriptive study [Lamb et al.], 371
- Bullerwell ML: methylene blue as an adjunct to treat vasoplegia in patients undergoing cardiac surgery requiring cardiopulmonary bypass: a literature review [Arevalo and Bullerwell], 455
- Bunger R: effect of norepinephrine on intracellular Ca²⁺ levels in malignant hyperthermia-susceptible B cells: pilot study in the search for a new diagnostic test for malignant hyperthermia [Perry et al.], 383
- Bupivacaine: implementation of education program for ultrasound-guided liposomal transverse abdominis plane (TAP) block protocol for open abdominal procedures, 479; minimum effective dose of intrathecal hyperbaric, for Cesarean delivery, 348
- ## C
- Calcium ion (Ca²⁺): effect of norepinephrine on intracellular levels in malignant hyperthermia, 383
- Calendar of Events, 81, 169, 249, 328, 417, 455
- Campbell Y: intravenous fat emulsion for treatment of local anesthetic systemic toxicity: best practice and review of the literature [Nedialkov et al.], 290
- Campero L: happy puppet syndrome: a case report of anesthetic management, 67
- Canale ML: implementation of a standardized handoff of anesthetized patients, 137
- Capacchione J: should continuous subcutaneous insulin infusion (CSII) pumps be used during the perioperative period? development of a clinical decision algorithm [Rotruck et al.], 194
- Cardiac surgery: methylene blue as adjunct in treating vasoplegia in patients requiring cardiopulmonary bypass, 455
- Cardiopulmonary bypass: as adjunct to treating vasoplegia in patients undergoing cardiac surgery, 455
- Case management: herbal medications and anesthesia, 242
- Case reports: of anesthetic management for happy puppet syndrome, 67; anesthetic management of a patient with type I Von Willebrand disease and uterine placental abruption, 209; anesthetic management of double-outlet right ventricle with Glenn shunt for Cesarean delivery, 408; complications related to complications related to laryngeal mask airway use and diagnosis of reinke edema, 443; emergent bilateral face-lift hematoma, 188; isolated harlequin syndrome following brachial plexus nerve block via interscalene approach, 56; point-of-care ultrasound gastric examination in management of ASA Class 3E patient, 379
- Ceremuga TE: investigation of the anxiolytic and antidepressant effects of Crocin, a compound from Saffron (*Crocus sativus* L), in the male Sprague-Dawley rat [Ceremuga et al.], 225
- Certified Registered Nurse Anesthetists (CRNA): adverse events during cosmetic surgery, 127; content and thematic analysis of closed claims resulting in death, 311; converging or diverging concepts in patient management and decision making, online; evidence-based use of nonopioid analgesics, 321; mobile computing devices in the perioperative environment and, 471; perceived knowledge and attitudes of, on fire risk assessment, 99; reference resource usage among nurse anesthetists, 299; transition to manager, 447; understanding motivations and barriers of, when involved in global health, 371
- Cesarean delivery: anesthetic management of double-outlet right ventricle with Glenn shunt for, 408; minimum effective dose intrathecal hyperbaric bupivacaine for, 348; pharmacologic methods for preventing pruritus in patients receiving intrathecal opioids for, 59; spinal-epidural anesthesia for, 109
- Checklists: evidence-based best practices of, in improving a regional training courses, 119
- Chicoine RW: investigation of the anxiolytic and antidepressant effects of Crocin, a compound from Saffron (*Crocus sativus* L), in the male Sprague-Dawley rat [Ceremuga et al.], 225
- Chinese herbal medicine: anesthesia and, 242
- Choi KH: effects of subanesthetic intravenous ketamine infusion on corticosterone and brain-derived neurotrophic factor in the plasma of male Sprague-Dawley rats [Radford et al.], 393
- Chun SM: investigation of the anxiolytic and antidepressant effects of Crocin, a compound from Saffron (*Crocus sativus* L), in the male Sprague-Dawley rat [Ceremuga et al.], 225
- Cigarette smoking: complications related to complications related to laryngeal mask airway use and diagnosis of reinke edema, 443
- Clayton BA: a content and thematic analysis of closed claims resulting in death [Hranchook et al.], 311; thematic analysis of the obstetric anesthesia cases from the AANA Foundation closed claims databank [Clayton et al.], 464; transfer of care in perioperative settings: a descriptive qualitative study [Wilbanks et al.], 401
- Clinical decision algorithm: use of continuous subcutaneous insulin infusion pumps during perioperative period, 194
- Clinical resources: usage among nurse anesthetists, 299
- Clinical technology: implementation of acceleromyography in increasing use of quantitative neuromuscular blockade monitoring, 269
- Closed claims: content and thematic analysis of, 311; identifying patterns across the AANA Foundation using thematic analysis, 27; improving patient outcomes through analysis of, 201; thematic analysis of adverse events during cosmetic surgery, 127; thematic analysis of obstetric anesthesia cases from the AANA Foundation closed claims databank, 464; transfer of care in perioperative settings: a descriptive qualitative study [Wilbanks et al.], 401
- Cobb C: understanding motivations and barriers of CRNAs involved in global health: a qualitative descriptive study [Lamb et al.], 371
- Coletto K: perceived knowledge and attitudes of certified

registered nurse anesthetists and student registered nurse anesthetists on fire risk assessment during time-out in the operating room [Coletto et al.], 99

Collaboration: team communication in the operating room, 11

Collins S: nitrous oxide for the management of labor analgesia [Collins et al.], 72

Communication: improved anesthesia handoff after implementation of the Written Handoff Anesthesia Tool, 361; team, in the operating room, 11

Complications. See also Postoperative complications: pharmacologic methods for preventing pruritus in patients receiving intrathecal opioids for Cesarean delivery, 59

Content analysis: of closed claims resulting in death, 311

Continuous subcutaneous insulin infusion pumps: use of during perioperative period, 194

Core temperature: relationship of tympanic and temporal temperature modalities to, in pediatric surgical patients, 19

Corticosterone: effects of subanesthetic intravenous ketamine infusion on BDNF in male Sprague-Dawley rats, 393

Cosmetic surgery: adverse events during, 127

Couture D: high rate of undiagnosed obstructive sleep apnea in patients undergoing total joint arthroplasty [Spence et al.], 282

Crawforth K: a content and thematic analysis of closed claims resulting in death [Hranchook et al.], 311

Credentialing: reference resource usage among nurse anesthetists, 299

Crerar C: implementation of an education program for an ultrasound-guided liposomal bupivacaine transversus abdominis plane (TAP) block protocol for open abdominal procedures [O'Driscoll et al.], 479

Crocin: investigation of the anxiolytic and antidepressant effects of, 225

Crosson JA: improve patient safety! recognizing and preventing residual neuromuscular blockade [Golinski et al], Supplement

Cryoneurolysis: cryoneurolysis for treating sensory nerve pain, 495

Cryoneurolysis: for treatment of sensory nerve pain, 495

Cuevas DK: complications related to laryngeal mask airway use and a postoperative diagnosis of reinke edema: a case study [Cuevas et al.], 443

D

Datta PK: anesthetic management of a unique case of double-outlet right ventricle with Glenn shunt for Cesarean delivery: a case report [Aravindan et al.], 408

Death: content and thematic analysis of closed claims resulting in, 311

Debriefing: evidence-based best practices of, in improving a regional training courses, 119

Decision making: converging or diverging concepts in patient management and, online

DeGroot JM: investigation of the anxiolytic and antidepressant effects of Crocin, a compound from Saffron (*Crocus sativus* L), in the male Sprague-Dawley rat [Ceremuga et al.], 225

Deliberate practice: evidence-based best practices of, in improving a regional training courses, 119

Descriptive studies: transfer of care in perioperative settings, 401; understanding motivations and barriers of CRNAs involved in global health, 371

Dexamethasone: intravenous, as an analgesic, 488

Diabetes mellitus: use of continuous subcutaneous insulin infusion pumps during perioperative period, 194

Diagnostic assay: pilot study in the search for new for

malignant hyperthermia, 383

Distractions: computing devices in the perioperative environment and, 471

Donoghue TJ: herbal medications and anesthesia case management, 242

Drug shortages: of propofol associated with postoperative nausea and vomiting outcomes, 147

Dunworth BA: implementation of acceleromyography to increase use of quantitative neuromuscular blockade monitoring: a quality improvement project [Dunworth et al.], 269

E

The Editor's Desk, 431; acetaminophen: oral or IV, does it matter?, 345; addicted to gaming? who says it is the real deal!, 266; aprotinin making a comeback? maybe, maybe not!, 265; association between wait time and 30-day mortality in adults undergoing hip fracture surgery, 185; blood donation by gay men under reconsideration, 9; breastfeeding and epidural analgesia with fentanyl, 10; cancer researcher at the Ohio State University resigns following multiple misconduct findings, 186; cloning monkeys in China, 97; color to an x-ray image coming your way?, 345; cows on Mars? not exactly, but there is a macromolecule a bit like methane!, 265; dysfunction of NaV1.4, a skeletal muscle voltage gated channel in sudden infant death syndrome, 185; fires in the OR, 10; fountain of youth in a mole rat, 98; the fountain of youth in a mole rat?, 98; frailty and cognitive impairment—a bad combination, 97; *global food security and wellness*, 186; hikers and campers beware: giardiasis can make you miserable!, 98; inefficiency in accessing clinical and research information, 346; liquid biopsy: cancer DNA in the blood?, 266; the LMA: development and preliminary trials of a new type of airway, 185; lowering the blood alcohol limit for driving?, 97; music therapy in the postop period, 97; the passing of Koko, 345; prevalence of needlestick injuries, attitude changes over 12 years in an urban academic hospital surgery department, 187; research involves very dangerous organisms, 9; science and clinical potpourri for your life and your practice, 97; structure-based discovery opioid analgesics with reduced side effects, 186; sugamadex and anaphylaxis: The Japanese experience, 266; typhoid fever: a dangerous, drug-resistant resurgence, 346; US in the lead: the supercomputing race is on! but for how long?, 265; what birds do for us, 10; who should take statins?, 9; worrying constructively?, 265

Education. See also AANA Journal Courses; Training courses: for an ultrasound-guided liposomal bupivacaine transversus abdominis plane (TAP) block protocol for open abdominal procedures, 479

Elective surgery: opioid-sparing effects of transversus abdominis plane block in hysterectomy, 41

Electronic medical record: implementation of acceleromyography in increasing use of quantitative neuromuscular blockade monitoring, 269

Emergence delirium: ketamine in decreasing, in the surgical patient with posttraumatic stress disorder, 220

Enhanced Recovery (ERAS) protocol: evidence-based use of nonopioid analgesics, 321; for patients undergoing total knee and total hip arthroplasty, 32

Epidural anesthesia: for double-outlet right ventricle with Glenn shunt for Cesarean delivery, 408

Epidural volume extension: effect of, on quality of combined spinal-epidural anesthesia for Cesarean delivery, 109

Errors. See Medical errors

Evidence based: best practices of simulation, checklists, deliberate practice and debriefing in developing a regional training course, 119; recommendations in improving patient outcomes with residual neuromuscular blockade, 157; reference resource usage among nurse anesthetists, 299; use of nonopioid analgesics, 321

Exparel: for an ultrasound-guided liposomal bupivacaine transversus abdominis plane (TAP) block protocol for open abdominal procedures, 479

Experts: converging or diverging concepts in patient management and decision making, online

F

Face-lift: facial hematoma as complication of, 188

Falyar CR: clinical application of point-of-care ultrasound gastric examination in the management of an ASA class 3E patient: a case report [Falyar and Kantzavelos], 379

Fasting times: point-of-care gastric ultrasound examination in management of ASA class 3E patient, 379

Fear conditioning: effects of subanesthetic intravenous ketamine infusion on corticosterone and BDNF in plasma of Sprague-Dawley rats, 393

Ferris M: reference resource usage among nurse anesthetists: a survey [Ferris & Muckle], 299

Fetal outcomes: with early neuraxial engagement in obese parturients, 234

Filler L: minimum effective dose (ED₅₀ and ED₉₅) of intrathecal hyperbaric bupivacaine for Cesarean delivery: a systematic review [Tubog et al.], 348

Fiore AT: nitrous oxide for the management of labor analgesia [Collins et al.], 72

Fire risk assessment: during time-out in the operating room, 99

First-time managers: transition of a Certified Registered Nurse Anesthetist's to, 447

Flanagan B: *AANA Journal* course–residual neuromuscular blockade: evidence-based recommendations to improve patient outcomes [Wiatrowski et al.], 157

Freeman K: *AANA Journal* course–residual neuromuscular blockade: evidence-based recommendations to improve patient outcomes [Wiatrowski et al.], 157

G

Garvin JT: reexamining metoclopramide's role in the prevention of postoperative nausea and vomiting: a secondary analysis [Masiongale et al.], 213

Gastric ultrasound: point-of-care examination in management of ASA class 3E patient, 379

Gastroesophageal reflux disease: complications related to complications related to laryngeal mask airway use and diagnosis of reinke edema, 443

Geisz-Everson M: a content and thematic analysis of closed claims resulting in death [Hranchook et al.], 311; transfer of care in perioperative settings: a descriptive qualitative study [Wilbanks et al.], 401

Geisz-Everson MA: thematic analysis of the obstetric anesthesia cases from the AANA Foundation closed claims databank [Clayton et al.], 464

Girolamo RG: implementation of an education program for an ultrasound-guided liposomal bupivacaine transversus abdominis plane (TAP) block protocol for open abdominal procedures [O'Driscoll et al.], 479

Glenn shunt: anesthetic management of double-outlet right ventricle with, for Cesarean delivery, 408

Global health: understanding motivations and barriers of CRNAs involved in, 371

Glutamate: ketamine in decreasing emergence delirium in, with posttraumatic stress disorder and, 220

Golinski M: adverse events during cosmetic surgery: a thematic analysis of closed claims [Golinski and Hranchook], 127; identifying patterns and meanings across the AANA Foundation closed claim dataset using thematic analysis methods, 27; improve patient safety! recognizing and preventing residual neuromuscular blockade [Golinski et al], Supplement

Gonzales C: understanding motivations and barriers of CRNAs involved in global health: a qualitative descriptive study [Lamb et al.], 371

Griffis CA: evidence-based use of nonopioid analgesics [Kremer & Griffis], 321

H

Handoff: implementation of standardized, of anesthetized patients, 137; improved anesthesia, after implementation of the Written Handoff Anesthesia Tool, 361

Han T: high rate of undiagnosed obstructive sleep apnea in patients undergoing total joint arthroplasty [Spence et al.], 282

Happy puppet syndrome: anesthetic management for, 67

Harenberg JL: opioid-sparing effects of transversus abdominis plane block in elective hysterectomy: a systematic review and meta-analysis [Tubog et al.], 41

Harlequin syndrome: isolated, following brachial plexus nerve block via interscalene approach, 56

Hematoma: emergent bilateral face-lift, 188

Henson DT: investigation of the anxiolytic and antidepressant effects of Crocin, a compound from Saffron (*Crocus sativus* L), in the male Sprague-Dawley rat [Ceremuga et al.], 225

Herbal medications: anesthesia and, 242

Herb-drug interactions: case management, 242

Hewer I: nitrous oxide for the management of labor analgesia [Collins et al.], 72

Hines CB: understanding bone cement implantation syndrome, 433

Hinson S: Hippocrates to Nightingale: converging or diverging concepts in patient management and decision making [Valdes et al.], online

Hintzsche K: subanesthetic-dose ketamine to decrease emergence delirium in the surgical patient with posttraumatic stress disorder, 220

Hirsch M: a content and thematic analysis of closed claims resulting in death [Hranchook et al.], 311

Hirsch MT: improving patient outcomes through closed-claims analysis: salient characteristics and patterns associated with respiratory events [Larson et al.], 201

Hooper VD: reexamining metoclopramide's role in the prevention of postoperative nausea and vomiting: a secondary analysis [Masiongale et al.], 213

Hranchook AM: adverse events during cosmetic surgery: a thematic analysis of closed claims [Golinski and Hranchook], 127; a content and thematic analysis of closed claims resulting in death [Hranchook et al.], 311; mobile computing devices in the perioperative environment: a survey exploring uses and experiences among certified registered nurse anesthetists [Hranchook et al.], 471

Humanitarianism: understanding motivations and barriers of CRNAs, when involved in global health, 371

Hybrid managers: transition of a Certified Registered Nurse Anesthetist's to, 447

Hyperglycemia: use of continuous subcutaneous insulin infusion pumps during perioperative period, 194

Hyperthermia: pilot study in the search for new diagnostic test for, 383
Hypoglossal nerve stimulation: as therapy for obstructive sleep apnea, 412
Hypoglycemia: use of continuous subcutaneous insulin infusion pumps during perioperative period, 194
Hypotension: methylene blue as adjunct in treating vasoplegia in patients undergoing cardiac surgery requiring cardiopulmonary bypass, 455; minimum effective dose of hyperbaric bupivacaine, for Cesarean delivery, 348
Hypoventilation: improving patient outcomes through closed-claims analysis, 201
Hysterectomy: opioid-sparing effects of transversus abdominis plane block in elective, 41

I

Informatics: computing devices in the perioperative environment, 471
Insulin: continuous subcutaneous infusion pumps during perioperative period, 194
Interdisciplinary communication: improving, in promoting patient safety, Supplement
Internet: reference resource usage among nurse anesthetists, 299
Interscalene approach: isolated harlequin syndrome following brachial plexus nerve block via, 56
Intravenous fat emulsion: for treatment of local anesthetic systemic toxicity, 290

J

Jordan L: a content and thematic analysis of closed claims resulting in death [Hranchook et al.], 311; improving patient outcomes through closed-claims analysis: salient characteristics and patterns associated with respiratory events [Larson et al.], 201

K

Kane J: effect of epidural volume extension on quality of combined spinal-epidural anesthesia for Cesarean delivery: a systematic review and meta-analysis [Kane et al.], 109
Kane T: effect of epidural volume extension on quality of combined spinal-epidural anesthesia for Cesarean delivery: a systematic review and meta-analysis [Kane et al.], 109
Kane TD: opioid-sparing effects of transversus abdominis plane block in elective hysterectomy: a systematic review and meta-analysis [Tubog et al.], 41
Kantzavelos L: clinical application of point-of-care ultrasound gastric examination in the management of an ASA class 3E patient: a case report [Falyar and Kantzavelos], 379
Kapanke K: perceived knowledge and attitudes of certified registered nurse anesthetists and student registered nurse anesthetists on fire risk assessment during time-out in the operating room [Coletto et al.], 99
Kasper CE: effect of norepinephrine on intracellular Ca^{2+} levels in malignant hyperthermia-susceptible B cells: pilot study in the search for a new diagnostic test for malignant hyperthermia [Perry et al.], 383
Keele R: relationship of Tympanic and Temporal Temperature Modalities to Core Temperature in Pediatric Surgical Patients [Minzola and Keele], 19
Ketamine: effects of subanesthetic intravenous on corticosterone and BDNF in the plasma of male Sprague-Dawley rats, 393; subanesthetic-dose, to decrease emergence delirium in the surgical patient with posttraumatic stress disorder, 220
Kirschbaum K: team communication in the operating room: a

measure of latent factors from a national sample of nurse anesthetists [Kirschbaum et al.], 11
Kradel B: Hippocrates to Nightingale: converging or diverging concepts in patient management and decision making [Valdes et al.], online
Kremer M: a content and thematic analysis of closed claims resulting in death [Hranchook et al.], 311
Kremer MJ: evidence-based use of nonopioid analgesics [Kremer & Griffis], 321
Kundu R: anesthetic management of a unique case of double-outlet right ventricle with glenn shunt for Cesarean delivery: a case report [Aravindan et al.], 408

L

Labor: nitrous oxide for the management of analgesia in, 72
Lambert LH: improved anesthesia handoff after implementation of the Written Handoff Anesthesia Tool (WHAT) [Lambert and Adams], 361
Lamb K: understanding motivations and barriers of CRNAs involved in global health: a qualitative descriptive study [Lamb et al.], 371
Larson SL: improving patient outcomes through closed-claims analysis: salient characteristics and patterns associated with respiratory events [Larson et al.], 201
Laryngeal mask airway use: complications related to and diagnosis of reinke edema, 443
Latent variables: in team communication in the operating room, 11
Lee Y-M: perceived knowledge and attitudes of certified registered nurse anesthetists and student registered nurse anesthetists on fire risk assessment during time-out in the operating room [Coletto et al.], 99
Lipid emulsion: intravenous fat emulsion for treatment of local anesthetic systemic toxicity, 290
Lipid rescue: intravenous fat emulsion for treatment of local anesthetic systemic toxicity, 290
Lipid sink: intravenous fat emulsion for treatment of local anesthetic systemic toxicity, 290
Liposomal bupivacaine: education for an ultrasound-guided transversus abdominis plane (TAP) block protocol for an open abdominal procedure, 479
Literature review: implementation of standardized handoff of anesthetized patients, 137; intravenous dexamethasone as an analgesic, 488; intravenous fat emulsion for treatment of local anesthetic systemic toxicity, 290; methylene blue as adjunct in treating vasoplegia in patients undergoing cardiac surgery requiring cardiopulmonary bypass, 455; reference resource usage among nurse anesthetists, 299
Local anesthetic systemic toxicity: intravenous fat emulsion for treatment of, 290
Looney SW: reexamining metoclopramide's role in the prevention of postoperative nausea and vomiting: a secondary analysis [Masiongale et al.], 213
Low dose: minimum effective dose of intrathecal hyperbaric bupivacaine, for Cesarean delivery, 348
Lutz C: implementation of acceleromyography to increase use of quantitative neuromuscular blockade monitoring: a quality improvement project [Dunworth et al.], 269; using evidence-based best practices of simulation, checklists, deliberate practice, and debriefing to develop and improve a regional anesthesia training course [Wiggins et al.], 119

M

Malignant hyperthermia: pilot study in the search for new diagnostic test for, 383

- Malpractice: adverse events during cosmetic surgery, 127; content and thematic analysis of closed claims resulting in death, 311; transfer of care in perioperative settings, 401
- Maresch KJ: *AANA Journal* Course: hypoglossal nerve stimulation: effective long-term therapy for obstructive sleep apnea, 412
- Martens J: a certified registered nurse anesthetist's transition to manager [Martens et al.], 447
- Martini L: *AANA Journal* course—residual neuromuscular blockade: evidence-based recommendations to improve patient outcomes [Wiatrowski et al.], 157
- Masiongale AJ: reexamining metoclopramide's role in the prevention of postoperative nausea and vomiting: a secondary analysis [Masiongale et al.], 213
- Masiongale J I: reexamining metoclopramide's role in the prevention of postoperative nausea and vomiting: a secondary analysis [Masiongale et al.], 213
- Mason-Nguyen J: opioid-sparing effects of transversus abdominis plane block in elective hysterectomy: a systematic review and meta-analysis [Tubog et al.], 41
- Maternal health: with early neuraxial engagement in obese parturients, 234; thematic analysis of obstetric anesthesia cases from the AANA Foundation closed claims databank, 464
- Matthews RW: a content and thematic analysis of closed claims resulting in death [Hranchook et al.], 311; improving patient outcomes through closed-claims analysis: salient characteristics and patterns associated with respiratory events [Larson et al.], 201
- McAuliffe MS: team communication in the operating room: a measure of latent factors from a national sample of nurse anesthetists [Kirschbaum et al.], 11
- McElhone P: a content and thematic analysis of closed claims resulting in death [Hranchook et al.], 311
- Medical errors: team communication in the operating room and, 11; thematic analysis of obstetric anesthesia cases from the AANA Foundation closed claims databank, 464
- Medical missions: understanding motivations and barriers of CRNAs, when involved in global health, 371
- Meta-analysis: effect of epidural volume extension on quality of combined spinal-epidural anesthesia for Cesarean delivery, 109; opioid-sparing effects of transversus abdominis plane block in elective hysterectomy, 41
- Methylene blue: as adjunct to treating vasoplegia in patients undergoing cardiac surgery, 455
- Methyl methacrylate: bone cement implementation syndrome and, 433
- Metoclopramide: role in the prevention of postoperative nausea and vomiting, 213
- Michaelson LP: effect of norepinephrine on intracellular Ca^{2+} levels in malignant hyperthermia-susceptible B cells: pilot study in the search for a new diagnostic test for malignant hyperthermia [Perry et al.], 383
- Minimum effective dose (Ed_{50} and Ed_{95}): of intrathecal hyperbaric bupivacaine for Cesarean delivery, 348
- Minzola DJ: isolated harlequin syndrome following brachial plexus nerve block via interscalene approach: a case report [Adams and Minzola], 56; relationship of Tympanic and Temporal Temperature Modalities to Core Temperature in Pediatric Surgical Patients [Minzola and Keele], 19
- Mitigation strategy: propofol drug shortage associated with worse postoperative nausea and vomiting outcomes despite, 147
- Mobile computing devices: in the perioperative environment, 471
- Moore C: implementation of an education program for an ultrasound-guided liposomal bupivacaine transversus abdominis plane (TAP) block protocol for open abdominal procedures [O'Driscoll et al.], 479
- Moore SG: intravenous dexamethasone as an analgesic: a literature review, 488
- Morbidity: thematic analysis of obstetric anesthesia cases from the AANA Foundation closed claims databank, 464
- Morphine consumption: in elective hysterectomy, 41
- Morrison S: implementation of acceleromyography to increase use of quantitative neuromuscular blockade monitoring: a quality improvement project [Dunworth et al.], 269; using evidence-based best practices of simulation, checklists, deliberate practice, and debriefing to develop and improve a regional anesthesia training course [Wiggins et al.], 119
- Morrison T: high rate of undiagnosed obstructive sleep apnea in patients undergoing total joint arthroplasty [Spence et al.], 282
- Mortality: thematic analysis of obstetric anesthesia cases from the AANA Foundation closed claims databank, 464
- Motz J: a certified registered nurse anesthetist's transition to manager [Martens et al.], 447
- Muckle TJ: reference resource usage among nurse anesthetists: a survey [Ferris & Muckle], 299
- Muldoon S: effect of norepinephrine on intracellular Ca^{2+} levels in malignant hyperthermia-susceptible B cells: pilot study in the search for a new diagnostic test for malignant hyperthermia [Perry et al.], 383
- Multimodal analgesia: intravenous dexamethasone as an analgesic, 488
- Murphy MJ: reexamining metoclopramide's role in the prevention of postoperative nausea and vomiting: a secondary analysis [Masiongale et al.], 213

N

- National Board of Certification and Recertification for Nurse Anesthetists: reference resource usage among nurse anesthetists, 299
- Naughton N: propofol drug shortage associated with worse postoperative nausea and vomiting outcomes despite a mitigation strategy [Neff et al.], 147
- Nausea and vomiting: propofol drug shortage associated with postoperative, 147; role of metoclopramide in the prevention of postoperative, 213
- Nedialkov AM: intravenous fat emulsion for treatment of local anesthetic systemic toxicity: best practice and review of the literature [Nedialkov et al.], 290
- Neff MP: propofol drug shortage associated with worse postoperative nausea and vomiting outcomes despite a mitigation strategy [Neff et al.], 147
- Neuraxial engagement: maternal and fetal outcomes with early, in obese parturients, 234
- Neurogenetic disorders: anesthetic management for happy puppet syndrome, 67
- Neuromuscular blockade: implementation of acceleromyography in increasing use of quantitative monitoring of, 269
- Nicely KW: a content and thematic analysis of closed claims resulting in death [Hranchook et al.], 311
- Nitrous oxide (N_2O): for the management of labor analgesia, 72
- Nonopioid analgesics: evidence-based use of, 321
- Norepinephrine: effect on intracellular Ca^{2+} levels in malignant hyperthermia, 383
- Novices: converging or diverging concepts in patient management and decision making, online
- Nurse anesthesia: identifying patterns across the AANA Foundation closed claim dataset using thematic analysis, 27
- Nurse anesthetists: reference resource usage among, 299;

on team communication in the operating room, 11;
understanding motivations and barriers of CRNAs, when
involved in global health, 371
Nurse managers: transition of a Certified Registered Nurse
Anesthetist's to, 447

O

Obesity: maternal and fetal outcomes with early neuraxial
engagement in parturients, 234
Obstetrics: anesthetic management of double-outlet right
ventricle with Glenn shunt for cesarean delivery,
408; minimum effective dose intrathecal hyperbaric
bupivacaine for Cesarean delivery, 348; nitrous oxide for
the management of labor analgesia, 72; pharmacologic
methods for preventing pruritus in patients receiving
intrathecal opioids for Cesarean delivery, 59; spinal-
epidural anesthesia for Cesarean delivery, 109; thematic
analysis of anesthesia cases from the AANA Foundation
closed claims databank, 464
Obstructive sleep apnea: long-term therapy for, 412;
undiagnosed, in patients undergoing total joint
arthroplasty, 282
Odom-Forren J: reexamining metoclopramide's role in the
prevention of postoperative nausea and vomiting: a
secondary analysis [Masiongale et al.], 213
O'Donnell J: using evidence-based best practices of
simulation, checklists, deliberate practice, and debriefing
to develop and improve a regional anesthesia training
course [Wiggins et al.], 119
O'Donnell JM: implementation of acceleromyography to
increase use of quantitative neuromuscular blockade
monitoring: a quality improvement project [Dunworth et
al.], 269
O'Driscoll L: implementation of an education program for
an ultrasound-guided liposomal bupivacaine transversus
abdominis plane (TAP) block protocol for open abdominal
procedures [O'Driscoll et al.], 479
Operating room: perceived knowledge and attitudes of CRNA
and SRNA on fire risk assessment during time-out in, 99;
team communication in the, 11
Opioids: pharmacologic methods for preventing pruritus in
patients receiving intrathecal, for Cesarean delivery, 59
Opioid-sparing effects: of transversus abdominis plane block in
elective hysterectomy, 41
Osborne-Smith L: effects of subanesthetic intravenous
ketamine infusion on corticosterone and brain-derived
neurotrophic factor in the plasma of male Sprague-Dawley
rats [Radford et al.], 393
Oseka L: anesthetic management in early recovery after
surgery protocols for total knee and total hip arthroplasty
[Oseka and Pecka], 32

P

Pain management: cryoneurolysis for treatment of sensory
nerve pain, 495
Pain scores: opioid-sparing effects of transversus abdominis
plane block in elective hysterectomy, 41
Park TY: effects of subanesthetic intravenous ketamine
infusion on corticosterone and brain-derived neurotrophic
factor in the plasma of male Sprague-Dawley rats [Radford
et al.], 393
Parturients: maternal and fetal outcomes with early neuraxial
engagement in obese, 234
Patient management: apnea in patients undergoing total joint
arthroplasty, 282; converging or diverging concepts in, online

Patient outcomes: improving through closed-claims analysis,
201; residual neuromuscular blockade, 157
Patient safety: fire risk assessment during time out in the
operating room, 99; identifying patterns across the
AANA Foundation closed claim dataset using thematic
analysis, 27; implementation of standardized handoff
of anesthetized patients, 137; improved anesthesia
handoff after implementation of the Written Handoff
Anesthesia Tool, 361; recognizing and preventing residual
neuromuscular blockade, Supplement
Pecka S: anesthetic management in early recovery after
surgery protocols for total knee and total hip arthroplasty
[Oseka and Pecka], 32
Pediatric surgical patients: relationship of tympanic and
temporal temperature modalities to core temperature in, 19
Penprase BB: mobile computing devices in the perioperative
environment: a survey exploring uses and experiences
among certified registered nurse anesthetists [Hranchook
et al.], 471
Perioperative settings: implementation of standardized
handoff of anesthetized patients, 137; mobile computing
devices in the, 471; transfer of care in, 401; use of
continuous subcutaneous insulin infusion pumps in, 194
Perry SM: effect of norepinephrine on intracellular Ca^{2+}
levels in malignant hyperthermia-susceptible B cells: pilot
study in the search for a new diagnostic test for malignant
hyperthermia [Perry et al.], 383
Pharmacologic methods: for preventing pruritus in patients
receiving intrathecal opioids for Cesarean delivery, 59
Phillips BJ: propofol drug shortage associated with worse
postoperative nausea and vomiting outcomes despite a
mitigation strategy [Neff et al.], 147
Physician manager: transition of a Certified Registered Nurse
Anesthetist's to, 447
Pilot studies: search for new diagnostic test for malignant
hyperthermia, 383
Piscotty RJ: mobile computing devices in the perioperative
environment: a survey exploring uses and experiences
among certified registered nurse anesthetists [Hranchook
et al.], 471
Placental abruption: anesthetic management of patient with
type 1 von Willebrand disease and abruption, 209
Polypoid degeneration: complications related to complications
related to laryngeal mask airway use and diagnosis of
reinke edema, 443
Postoperative complications: emergent bilateral face-lift
hematoma, 188; implementation of acceleromyography
in increasing use of quantitative neuromuscular blockade
monitoring, 269; metoclopramide's role in the prevention
of postoperative nausea and vomiting, 213
Postoperative diagnoses: complications related to laryngeal
mask airway use and, 443
Postoperative pain management: isolated harlequin syndrome
following brachial plexus nerve block via interscalene
approach, 56
Posttraumatic stress disorder (PTSD): ketamine in decreasing
emergence delirium in the surgical patient with, 220
Powel A: complications related to laryngeal mask airway use
and a postoperative diagnosis of reinke edema: a case
study [Cuevas et al.], 443
Prevention: thematic analysis of obstetric anesthesia cases for
the AANA Foundation closed claims databank, 464
Prophylactic antiemetics: propofol drug shortage associated
with postoperative nausea and vomiting, 147
Prophylaxis: metoclopramide's role in the prevention of
postoperative nausea and vomiting, 213

Propofol: drug shortage associated with worse postoperative nausea and vomiting outcomes, 147
Pruritus: pharmacologic methods for preventing, in patients receiving intrathecal opioids for Cesarean delivery, 59
Pugh MA: pharmacologic methods for preventing pruritus in patients receiving intrathecal opioids for Cesarean delivery [Vice-O'con et al.], 59
Pulmonary aspiration: in management of ASA class 3E patient, 379

Q

Quality management: effect of epidural volume extension on quality of combined spinal-epidural anesthesia for Cesarean delivery, 109; implementation of acceleromyography in increasing use of quantitative neuromuscular blockade monitoring, 269
Quantitative monitoring: implementation of acceleromyography in increasing use of quantitative neuromuscular blockade monitoring, 269
Quantitative neuromuscular blockade monitoring: implementation of acceleromyography in increasing use of, 269
Questionnaire: undiagnosed obstructive sleep apnea in patients undergoing total joint arthroplasty, 282

R

Radford KD: effects of subanesthetic intravenous ketamine infusion on corticosterone and brain-derived neurotrophic factor in the plasma of male Sprague-Dawley rats [Radford et al.], 393
Ramsey VL: minimum effective dose (ED₅₀ and ED₉₅) of intrathecal hyperbaric bupivacaine for Cesarean delivery: a systematic review [Tubog et al.], 348
Randall SA: investigation of the anxiolytic and antidepressant effects of Crocin, a compound from Saffron (*Crocus sativus* L), in the male Sprague-Dawley rat [Ceremuga et al.], 225
Rats. See Sprague-Dawley rats
Ray S: maternal and fetal outcomes with early neuraxial engagement in obese parturients [Ray and Bradley], 234
Reference materials: reference resource usage among nurse anesthetists, 299
Regional anesthesia: evidence-based best practices of simulation, checklists, deliberate practice and debriefing in developing a training course, 119; isolated harlequin syndrome following brachial plexus nerve block via interscalene approach, 56
Reinke edema: complications related to complications related to laryngeal mask airway use and diagnosis of, 443
Residual neuromuscular blockade: evidence-based recommendations in improving patient outcomes, 157; implementation of acceleromyography in increasing use of quantitative, 269; recognizing and preventing, Supplement
Respiratory care: complications related to laryngeal mask airway use and, 443
Respiratory events: improving patient outcomes through closed-claims analysis, 201
Reversal agents: in preventing residual neuromuscular blockade, Supplement
Rhytidectomy: facial hematoma as complication of, 188
Risk factors: with emergent bilateral face-lift hematoma, 188; propofol drug shortage associated with postoperative nausea and vomiting, 147
Risks: metoclopramide's role in the prevention of postoperative nausea and vomiting, 213
Roberson MC: anesthetic management of a patient with type 1 Von Willebrand disease and uterine placental abruption:

a case report [Roberson et al.], 209
Rotruck J: should continuous subcutaneous insulin infusion (CSII) pumps be used during the perioperative period? development of a clinical decision algorithm [Rotruck et al.], 194
Rotruck S: should continuous subcutaneous insulin infusion (CSII) pumps be used during the perioperative period? development of a clinical decision algorithm [Rotruck et al.], 194

S

Saffron: investigation of the anxiolytic and antidepressant effects of Crocin, a compound from, 225
Sandberg WS: implementation of acceleromyography to increase use of quantitative neuromuscular blockade monitoring: a quality improvement project [Dunworth et al.], 269
Secondary analysis: metoclopramide's role in the prevention of postoperative nausea and vomiting, 213
Sensory nerve pain: cryoneurolysis for treatment of, 495
Shanks A: propofol drug shortage associated with worse postoperative nausea and vomiting outcomes despite a mitigation strategy [Neff et al.], 147
Simulation: evidence-based best practices of, in improving a regional training courses, 119
Sloan N: AANA *Journal* course—residual neuromuscular blockade: evidence-based recommendations to improve patient outcomes [Wiatrowski et al.], 157
Smiley A: AANA *Journal* course—cryoneurolysis for the treatment of sensory nerve pain [Smiley and McGuire], 495
Spence D: high rate of undiagnosed obstructive sleep apnea in patients undergoing total joint arthroplasty [Spence et al.], 282
Spinal anesthesia: for Cesarean delivery, 109; minimum effective dose of hyperbaric bupivacaine, for Cesarean delivery, 348
Sprague-Dawley rats: effects of subanesthetic intravenous ketamine infusion on corticosterone and BDNF in plasma of, 393; investigation of the anxiolytic and antidepressant effects of Crocin, 225
Standard of care: improving patient outcomes through closed-claims analysis, 201
Stanley LR: investigation of the anxiolytic and antidepressant effects of Crocin, a compound from Saffron (*Crocus sativus* L), in the male Sprague-Dawley rat [Ceremuga et al.], 225
STOP-BANG Questionnaire: undiagnosed obstructive sleep apnea in patients undergoing total joint arthroplasty, 282
Student registered nurse anesthetists (SRNA): perceived knowledge and attitudes of, on fire risk assessment, 99
Stump L: a certified registered nurse anesthetist's transition to manager [Martens et al.], 447
Subramaniam R: anesthetic management of a unique case of double-outlet right ventricle with glenn shunt for Cesarean delivery: a case report [Aravindan et al.], 408
Surgical patients: ketamine in decreasing emergence delirium in, with posttraumatic stress disorder, 220; relationship of tympanic and temporal temperature modalities to core temperature in pediatric, 19
Surgical time outs: fire risk assessment during, in the operating room, 99
Suszan L: should continuous subcutaneous insulin infusion (CSII) pumps be used during the perioperative period? development of a clinical decision algorithm [Rotruck et al.], 194
Swanson M: team communication in the operating room: a measure of latent factors from a national sample of nurse

anesthetists [Kirschbaum et al.], 11
Sympathetic chain: isolated harlequin syndrome following brachial plexus nerve block via interscalene approach and, 56
Systematic review: effect of epidural volume extension on quality of combined spinal-epidural anesthesia for Cesarean delivery, 109; of intrathecal hyperbaric bupivacaine for Cesarean delivery, 348

T

Talawar P: anesthetic management of a unique case of double-outlet right ventricle with glenn shunt for Cesarean delivery: a case report [Aravindan et al.], 408
Tariman JD: perceived knowledge and attitudes of certified registered nurse anesthetists and student registered nurse anesthetists on fire risk assessment during time-out in the operating room [Coletto et al.], 99
TeamSTEPPS: implementation of a standardized handoff of anesthetized patients, 137
Teamwork: converging or diverging concepts in patient management and decision making, online; with nurse anesthetists in the operating room, 11
Temperature monitoring: relationship of tympanic and temporal temperature modalities to core, in pediatric surgical patients, 19
Temporal temperature: relationship to core temperature in pediatric surgical patients, 19
Thematic analysis: adverse events during cosmetic surgery, 127; of closed claims resulting in death, 311; identifying patterns across the AANA Foundation closed claim dataset using, 27; of obstetric anesthesia cases from the AANA closed claims databank, 464
Thompson A: propofol drug shortage associated with worse postoperative nausea and vomiting outcomes despite a mitigation strategy [Neff et al.], 147
Time outs: fire risk assessment during, in the operating room, 99
Todd LA: should continuous subcutaneous insulin infusion (CSII) pumps be used during the perioperative period? development of a clinical decision algorithm [Rotruck et al.], 194
Total joint arthroplasty: anesthesia management in early recovery after surgical protocols for, 32; undiagnosed obstructive sleep apnea in patients undergoing, 282
Training courses: evidence-based best practices of simulation, checklists, deliberate practice and debriefing in developing, 119
Transfer of care: in perioperative settings, 401
Transverse abdominis plane (TAP) block: education program for an ultrasound-guided liposomal bupivacaine protocol for open abdominal procedures, 479
Transversus abdominis plane (TAP) block: opioid-sparing effects of, in elective hysterectomy, 41
Tubog TD: effect of epidural volume extension on quality of combined spinal-epidural anesthesia for Cesarean delivery: a systematic review and meta-analysis [Kane et al.], 109; minimum effective dose (ED_{50} and ED_{95}) of intrathecal hyperbaric bupivacaine for Cesarean delivery: a systematic review [Tubog et al.], 348; opioid-sparing effects of transversus abdominis plane block in elective hysterectomy: a systematic review and meta-analysis [Tubog et al.], 41
Tympanic temperature: relationship to core temperature in pediatric surgical patients, 19

U

Ultrasound gastric examination: in management of ASA class 3E patient, 379

Umadhay T: intravenous fat emulsion for treatment of local anesthetic systemic toxicity: best practice and review of the literature [Nedialkov et al.], 290
Upper air stimulation: long-term therapy for obstructive sleep apnea, 412
Uterine placental abruption: anesthetic management of patient with type 1 von Willebrand disease and, 209

V

Valdes JA: intravenous fat emulsion for treatment of local anesthetic systemic toxicity: best practice and review of the literature [Nedialkov et al.], 290; Hippocrates to Nightingale: converging or diverging concepts in patient management and decision making [Valdes et al.], online
Vasoplegia: methylene blue as adjunct in treating patients undergoing cardiac surgery, 455
Vice-O'Con K: pharmacologic methods for preventing pruritus in patients receiving intrathecal opioids for Cesarean delivery [Vice-O'con et al.], 59
Vigersky R: should continuous subcutaneous insulin infusion (CSII) pumps be used during the perioperative period? development of a clinical decision algorithm [Rotruck et al.], 194
Vigilance: computing devices in the perioperative environment and, 471
Voepel-Lewis T: propofol drug shortage associated with worse postoperative nausea and vomiting outcomes despite a mitigation strategy [Neff et al.], 147
Voluntarism: understanding motivations and barriers of CRNAs, when involved in global health, 371
Vomiting. *See* Nausea and vomiting
von Willebrand disease: anesthetic management of a patient with type 1, and uterine placental abruption, 209

W

Wagner D: propofol drug shortage associated with worse postoperative nausea and vomiting outcomes despite a mitigation strategy [Neff et al.], 147
Wanderer JP: implementation of acceleromyography to increase use of quantitative neuromuscular blockade monitoring: a quality improvement project [Dunworth et al.], 269
Whitney A-M: emergent bilateral face-lift hematoma: a case report, 188
Wiatrowski R: AANA *Journal* course—residual neuromuscular blockade: evidence-based recommendations to improve patient outcomes [Wiatrowski et al.], 157
Wiggins LL: using evidence-based best practices of simulation, checklists, deliberate practice, and debriefing to develop and improve a regional anesthesia training course [Wiggins et al.], 119
Wigley MD: anesthetic management of a patient with type 1 Von Willebrand disease and uterine placental abruption: a case report [Roberson et al.], 209
Wilbanks B: thematic analysis of the obstetric anesthesia cases from the AANA Foundation closed claims databank [Clayton et al.], 464
Wilbanks BA: transfer of care in perioperative settings: a descriptive qualitative study [Wilbanks et al.], 401
Wilkins K: propofol drug shortage associated with worse postoperative nausea and vomiting outcomes despite a mitigation strategy [Neff et al.], 147
Williamson W: implementation of an education program for an ultrasound-guided liposomal bupivacaine transversus abdominis plane (TAP) block protocol for open abdominal

procedures [O'Driscoll et al.], 479
Wofford K: implementation of an education program for
an ultrasound-guided liposomal bupivacaine transversus
abdominis plane (TAP) block protocol for open abdominal
procedures [O'Driscoll et al.], 479

Wright SM: improve patient safety! recognizing and
preventing residual neuromuscular blockade [Golinski et
al], Supplement
Written Handoff Anesthesia Tool (WHAT): improved
anesthesia handoff after implementation of, 361